

The Dopamine Stabilizers (S)-(-)-(3-Methanesulfonyl-phenyl)-1-propyl-piperidine [(–)-OSU6162] and 4-(3-Methanesulfonyl-phenyl)-1-propyl-piperidine (ACR16) Show High in Vivo D₂ Receptor Occupancy, Antipsychotic-Like Efficacy, and Low Potential for Motor Side Effects in the Rat

Sridhar Natesan, Kjell A. Svensson, Greg E. Reckless, José N. Nobrega, Karen B. L. Barlow, Anette M. Johansson, and Shitij Kapur

Schizophrenia Program and PET Centre (S.N., G.E.R., S.K.) and Neuroimaging Research Section (J.N.N., K.B.L.B.), Centre for Addiction and Mental Health, Toronto, Ontario, Canada; Lilly Research Laboratories, Indianapolis, Indiana (K.A.S., A.M.J.); and Departments of Pharmacology (J.N.N.) and Psychiatry (S.K.), University of Toronto, Toronto, Ontario, Canada

Received February 13, 2006; accepted April 26, 2006

ABSTRACT

“Dopamine stabilizers” are a new class of compounds that have the ability to reverse both hypo- as well as hyperdopaminergia in vivo. This class, exemplified by the phenylpiperidines (S)-(-)-3-(3-methanesulfonyl-phenyl)-1-propyl-piperidine [(–)-OSU6162] and 4-(3-methanesulfonyl-phenyl)-1-propyl-piperidine [ACR16] although lacking high in vitro binding affinity for dopamine D₂ receptor [(–)-OSU6162, K_i = 447 nM; ACR16, K_i > 1 μM], shows functional actions, suggestive of their interaction. Hence, we evaluated in vivo D₂ occupancy of these agents in rats and correlated it to observed effects in a series of behavioral, neurochemical, and endocrine models relevant to the dopamine system and antipsychotic effect. Both (–)-OSU6162 and ACR16 showed robust dose-dependent striatal D₂ occupancy with ED₅₀ values of 5.27 and 18.99 mg/kg s.c., respectively, and functional assays showed no partial agonism. Over an occupancy range of 37 to 87% (3–60

mg/kg) for (–)-OSU6162 and 35 to 74% (10–60 mg/kg) for ACR16, we observed both inhibitory (amphetamine-induced locomotor activity) and stimulatory effects (in habituated rats). Haloperidol, over a similar occupancy range (33–78%), potently inhibited psychostimulant activity and induced catalepsy, but it failed to activate habituated animals. In the conditioned avoidance response assay, ACR16 was clearly more efficacious than (–)-OSU6162. In addition, both these compounds demonstrated significant preferential Fos induction in the nucleus accumbens compared with the dorsolateral striatum, a strong predictor of atypical antipsychotic efficacy. The results suggest that dopamine stabilizers exhibit locomotor stabilizing as well as antipsychotic-like effects, with low motor side effect liability, in a dose range that corresponds to high D₂ in vivo occupancy.

¹Dysregulation or disruption of dopaminergic function underlies a variety of neuropsychiatric disorders, including schizophrenia and Parkinson's disease. Schizophrenia is thought to result from subcortical hyperdopaminergia (leading to positive symptoms) and cortical hypodopaminergia (leading to negative and cognitive deficits) (Abi-Dargham, 2004). Efforts to treat symptoms of schizophrenia have focused on blocking dopaminergic activity, and although the introduction of antipsychotics has revolutionized the treatment of this debilitating disorder, there remains an unmet

need to treat negative symptoms and cognitive deficits (Barbeau, 1961; Seeman, 1987; Nieoullon and Coquerel, 2003; Carlsson et al., 2004). Thus, optimal treatment for these conditions may not rely solely on dopamine blockade but may also include a dopamine tone-dependent modulation or “stabilization” of the dopaminergic system. There is now a class of drugs that has been suggested to possess such properties, exemplified by (–)-OSU6162 and ACR16 (Sonnesson et al., 1994; Ekesbo et al., 1997; Tedroff et al., 1998; Nichols et al., 2002; Carlsson et al., 2004; Nilsson et al., 2004; Rung et al., 2005). These compounds are reported to lack high in vitro affinity for various neuroreceptors. For (–)-OSU6162, D₂ and D₃ K_i values were 447 and 1305, nM,

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.
doi:10.1124/jpet.106.102905.

ABBREVIATIONS: (–)-OSU6162, (S)-(-)-3-(3-methanesulfonyl-phenyl)-1-propyl-piperidine; ACR16, 4-(3-methanesulfonyl-phenyl)-1-propyl-piperidine; MK-801, 5*H*-dibenzo[*a,c*]cyclohept-5,10-imine (dizocilpine maleate); CAR, conditioned avoidance response; NSD1015 HCl, 3-hydroxybenzyl hydrazine hydrochloride; CI, confidence interval; ANOVA, analysis of variance Di-PR-5,6-ADTN, *N,N*-dipropylamino-5,6-dihydroxytetralin.

respectively, and were reported with K_i values $>1 \mu\text{M}$ for other targets (Sonesson et al., 1994), whereas for ACR16, the K_i values for a number of targets, including monoaminergic receptors, were reported to be $>10 \mu\text{M}$ (Pettersson et al., 2002). Similar results were found by researchers at Eli Lilly with D_2/D_3 K_i values of 23/2.2 and 0.9/2.8 μM for ACR16 and (-)-OSU6162, respectively, using 7-hydroxy-2-dipropylaminotetralin as radioligand in cloned cells transfected with human D_2/D_3 receptors (S. Little, personal communication).

(-)-OSU6162 (also called PNU-96391A) was identified among a series of substituted (*S*)-phenylpiperidines examined for their ability to interact with central dopamine receptors, and it was found to be highly active in vivo on the synthesis and turnover of dopamine (Sonesson et al., 1994). It also caused inhibition of amphetamine-induced hyperlocomotion and stimulation of locomotor activity in rats with low psychomotor activity without causing motor impairment or catalepsy (Svensson et al., 1993; Waters et al., 1993; Sonesson et al., 1994). This class of compounds has been named "dopamine stabilizers" based on the ability of these compounds to either stimulate, suppress, or show no effect on motor activity, depending upon the prevailing dopaminergic tone (Carlsson et al., 2004). In search for compounds through a similar in vivo testing strategy, a close structural analog, ACR16, emerged as a molecule displaying a similar pharmacological profile as that of (-)-OSU6162. It induced a dose-dependent increase in the dopamine metabolite 3,4-dihydroxyphenylacetic acid in the striatum, cortex, and limbic areas, whereas exploratory locomotor activity was unaffected. In addition, it increased locomotor activity in habituated animals but antagonized amphetamine-, MK-801-, and cocaine-induced locomotor activity in mice, and recently, it has been shown to reverse social withdrawal induced by MK-801 in rats (Carlsson, 2002; Waters et al., 2002; Nilsson et al., 2004).

Although these drugs show intriguing abilities to modify dopamine-related behaviors in a bidirectional manner, the mechanisms by which these effects are induced are not completely understood. Although these compounds show weak in vitro D_2 receptor affinities (see above), a positron emission tomography study in the rhesus monkey showed that continuous intravenous administration of (-)-OSU6162 produced nearly 80% displacement of [^{11}C]raclopride in the striatal brain area (Neu et al., 1997; Tedroff et al., 1998; Ekesbo et al., 1999). Thus, our first objective in this study was to evaluate the in vivo rat striatal dopamine D_2 receptor occupancy and catalepsy induction by (-)-OSU6162 and ACR16, over their effective dose ranges, using previously validated methods (Wadenberg et al., 2000). Second, we examined the ability of these compounds to stabilize motor function in animals made hyperactive by means of *d*-amphetamine pretreatment and also examined whether the compounds could affect motor activity in rats with low baseline activity as the result of repeated habituation to the activity cages. To examine whether stabilization could be obtained with a standard dopamine D_2 blocker, we included haloperidol at doses that caused equivalent D_2 occupancy as those induced by (-)-OSU6162 and ACR16. To further characterize the functional effects of these drugs, we examined whether ACR16 and (-)-OSU6162 would show partial agonist activity in vivo at the dopamine D_2 autoreceptors and also examined whether

they could increase plasma levels of prolactin. Finally, the abilities of these compounds to inhibit conditioned avoidance responding (CAR) (Wadenberg et al., 2000) and to induce expression of Fos in the nucleus accumbens (Deutch et al., 1992; Robertson et al., 1994) were used as additional assessments of their antipsychotic potential.

Materials and Methods

Animals. The experiments were carried out on adult male Sprague-Dawley rats weighing 250 to 275 g when procured from Charles River (Montreal, QC, Canada). They were housed two per cage on a 12:12 reverse light/dark cycle (lights off at 8:00 AM) with free access to food and water. The animals were allowed to acclimatize for a minimum of 5 days before being used for experimentation. All experiments were approved by the Institution's Animal Care Committee. For the reserpine study, male Sprague-Dawley rats ($n = 5$ –6/group) weighing 200 to 250 g were obtained from Harlan (Indianapolis, IN). The rats were acclimated for 1 week before testing. The study protocol was approved by the Animal Care and Use Committee of Eli Lilly & Co. (Indianapolis, IN).

Drugs. (-)-OSU6162 and ACR16 were synthesized at Eli Lilly & Co., whereas haloperidol was obtained from Sabex Inc. (Boucherville, QC, Canada). All drugs were dissolved in 1% glacial acetic acid and were administered s.c. in a volume of 1 ml/kg body weight. [^3H]raclopride (PerkinElmer Life and Analytical Sciences, Boston, MA) was used as the radioligand for the occupancy study and was administered i.v. via the tail vein. *d*-Amphetamine sulfate was obtained from U.S. Pharmacopoeia (Rockville, MD) and dissolved in physiological saline and administered subcutaneously in a volume of 1 ml/kg for the locomotor experiments. Reserpine (Sigma/RBI, Natick, MA) was dissolved in a few drops of glacial acetic acid and made up to volume in 5% glucose solution. The *l*-aromatic amino acid decarboxylase inhibitor 3-hydroxybenzylhydrazine hydrochloride (NSD1015 HCl; Sigma-Aldrich, St. Louis, MO) was dissolved in saline. Both reserpine and NSD1015 HCl were injected in a volume of 5 ml/kg subcutaneously.

Striatal Dopamine D_2 Receptor Occupancy Experiments. (-)-OSU6162 (3–120 mg/kg), ACR-16 (10–240 mg/kg), and haloperidol (0.025–1 mg/kg) were administered to rats to obtain a dose response of D_2 receptor occupancy levels, 1 h after drug administration. Animals ($n = 5$) were randomly assigned to each dose level. Thirty minutes before sacrifice, all animals received an intravenous injection of [^3H]raclopride [7.5 $\mu\text{Ci}/\text{rat}$, in a volume of 0.4 ml of 0.9% (w/v) NaCl solution] to determine D_2 occupancy. Animals were sacrificed by decapitation, and striata and cerebellum were rapidly dissected. Approximately one-third of the cerebellum and the left and right striata pooled as a single sample were collected. The tissue was dissolved in 2 ml of Solvable (Canberra Packard Canada, Montreal, QC, Canada) and was gently agitated for 24 h. Scintillation fluid was added, and the vials were allowed to shake for another 24 h. Radioactivity was determined using liquid scintillation spectrometry using an LS5000 CE liquid scintillation counting system (Beckman Coulter, Fullerton, CA). To obtain an index of the binding potential of the dopamine D_2 receptors, the ratio of striatum minus cerebellum (index of specific binding)/cerebellum (index of free and nonspecific binding), a clinical method validated for experimental animals, was used (Wadenberg et al., 2000). To determine D_2 occupancies as a function of time, 30 mg/kg (-)-OSU6162, 60 mg/kg ACR-16, and 0.5 mg/kg haloperidol were evaluated at different time points. The occupancy induced by the drug of interest was calculated using the formula: %Occupancy = $100 \times (D_2BP_{\text{controls}} - D_2BP_{\text{drug}} / D_2BP_{\text{controls}})$, where D_2BP_{controls} is the pooled D_2 binding potential of all the control animals, and D_2BP_{drug} is the D_2 binding potential of a drug-treated animal. Occupancy curves and the ED_{50} values (dose at which 50% receptors are occupied) were determined using the non-

linear regression equation representing a rectangular hyperbola [$y = ax/(b + y)$] using SigmaPlot (SPSS Inc., Chicago, IL).

Catalepsy. Animals used for the occupancy experiments were also used to measure catalepsy. Catalepsy was measured 10 min before sacrifice. Animals were placed on an inclined grid (60°), and the time the animals remained immobile (excluding the first 30 s) was used as an index of catalepsy (on a scale of 0–5 in which time was a square-root transformation: 0, 0 to 0.08; 1, 0.09 to 0.35; 2, 0.36 to 0.80; 3, 0.81 to 1.42; 4, 1.43 to 2.24; and 5, >2.24 min (Wadenberg et al., 2000). An animal with a score of 2 or greater was considered cataleptic.

Behavioral Stabilization—Locomotor Activity. Custom-made locomotor activity boxes similar to the home cages (clear Plexiglas, 27 × 48 × 20 cm), equipped with a row of six photocell beams placed 3 cm above the floor of the cage were used. A computer was used to record disruption of the photocell beams. Two sets of the experiment were carried out. In case of amphetamine-induced hyperactivity, 30 min after drug administration, rats were placed in the locomotor activity boxes to habituate for a period of 30 min. After 30 min, *d*-amphetamine (1.5 mg/kg s.c.) was administered, and locomotor activity was monitored for a period of 60 min. The ED₅₀ value was the dose that was required to inhibit 50% of locomotor activity counts, recorded over the period of 60 min, with respect to vehicle-treated amphetamine-administered animals and was calculated using nonlinear regression using SigmaPlot. In the case of locomotor activity in habituated rats, they were habituated to the locomotor boxes for three consecutive days for a period of 1 h each day. On the day of the experiment, rats were allowed to habituate to the locomotor activity cage for 30 min followed by an injection of the drug. Locomotor activity was monitored for a period of 1 h. Each dose of drug testing had a minimum of five animals. We tested 3 to 60 mg/kg (–)-OSU6162, 3 to 60 mg/kg ACR16, and 0.01 to 0.5 mg/kg haloperidol in amphetamine-induced hyperlocomotor activity, whereas 1 to 60 mg/kg (–)-OSU6162, 3 to 60 mg/kg ACR16, and 0.01 to 0.1 mg/kg haloperidol were tested in habituated rats.

Striatal DOPA Levels in Reserpinized Rats. Rats were first dosed with reserpine (5 mg/kg s.c.; 18–20 h before) to deplete their monoamine vesicular stores. Test compounds were given subcutaneously 30 min before the *L*-aromatic amino acid decarboxylase inhibitor NSD1015 HCl (100 mg/kg s.c.), and rats were sacrificed 30 min later. As a result of prolonged synaptic depletion of dopamine due to reserpine, the synthesis-regulating dopamine D₂ autoreceptors developed a certain degree of supersensitivity and are hence sensitive to dopamine agonists/partial agonists (Svensson et al., 1991). In the presence of an *L*-aromatic amino acid decarboxylase inhibitor, *L*-DOPA accumulates and the amount of *L*-DOPA accumulated during a 30-min period is taken as a measure of the synthesis rate of dopamine. Partial and full agonists in reserpinized rats decrease this endpoint. Each treatment group had a minimum of five animals. Striatal levels of *L*-DOPA were measured using standard high-pressure liquid chromatography with electrochemical detection techniques.

Conditioned Avoidance Response. Rats were trained and tested in a two-way active avoidance apparatus (custom made shuttle boxes; MED Associates, St. Albans, VT). The boxes contained a sound- and light-attenuating shell in which two compartments of equal size were separated by a translucent partition with a single opening to ensure a two-way active avoidance setting. The shuttle boxes were enabled with a tilting grid floor and microswitch detection. An 80-dB white noise served as a conditioned stimulus, whereas a 0.8-mA foot shock served as the unconditioned stimulus. The rat's location was detected by activation of microswitches fixed at the base of each compartment and programs running on a computer controlled the operations of the task. Animals that moved to the other side of the box within the period of the conditioned stimulus (10 s) were noted as having made an "avoidance" response. Those who escaped the shock in the next 20 s were termed as having "escaped," and those not escaping within the total 30 s were termed as "escape

failures" (Wadenberg et al., 2001). Rats were trained for 5 days before drug testing. A performance criterion of greater than 80% avoidance after the 5-day training served as the basis for selecting rats that were used for drug testing. The entire protocol as well as recording of the performance of the animal was controlled by MED Associates computer routines. (–)-OSU6162 (30, 60, and 120 mg/kg; *n* = 5), ACR16 (30, 60, and 120 mg/kg; *n* = 7), and haloperidol (0.01, 0.05, and 0.15 mg/kg; *n* = 6) were administered in a manner such that animals in each drug group served as their own controls in a within-subject design. CAR was measured at 0, 20, 90, and 240 min and finally at 24 h after drug administration, and an interval of 2 days served as a washout period.

Fos Immunohistochemistry. Rats were deeply anesthetized with sodium pentobarbital (100 mg/kg i.p.) 2 h after drug administration. They were perfused through the ascending aorta with saline followed by 4% paraformaldehyde in 0.1 M phosphate buffer. Whole brains were collected and postfixed in 4% paraformaldehyde for 24 h, transferred to sucrose solutions (10% for 2 h, 20% for 12 h, and 30% for 24 h), and then dried and stored at –80°C until processing. Free-floating, 40- μ m cryostat sections were incubated with a polyclonal primary antiserum raised in rabbit against the Fos peptide (4–17 amino acids of human Fos; Calbiochem, San Diego, CA), diluted 1:5000 for 48 h at 4°C. The tissue sections were then exposed to biotinylated goat anti-rabbit secondary antibody (1:200; Vector Laboratories, Burlingame, CA), which was followed by incubation with horseradish peroxidase avidin-biotin complex (Vector Laboratories) to visualize the Fos staining. Fos-immunoreactive nuclei were counted by an observer blind to treatment conditions, within a 400 × 400- μ m grid at a magnification of 100× in the shell of the nucleus accumbens and dorsolateral striatum (bregma, 1.70 to 1.00; Paxinos and Watson, 1986; Robertson et al., 1994) using an MCID M5 system (Imaging Research, St. Catharines, ON, Canada). Cell counts were obtained from at least three separate brain sections for each brain, obtained from four subjects per group. (–)-OSU6162 (10, 30, 60, and 120 mg/kg), ACR16 (10, 30, 60, and 120 mg/kg), and haloperidol (0.01, 0.05, 0.5, and 1 mg/kg) were tested for Fos protein expression.

Prolactin Estimation. Prolactin levels were measured using plasma collected from rats sacrificed for the D₂ receptor occupancy/catalepsy experiments. Plasma samples were stored at –80°C until they were assayed. Prolactin levels (nanograms per milliliter) were measured using a rat prolactin enzyme immunoassay kit (ALPCO Diagnostics, Windham, NH).

Results

Occupancy and Catalepsy. All three drugs showed dose-dependent striatal D₂ receptor occupancy (Fig. 1). (–)-OSU6162 over a range of 3 to 120 mg/kg (37–90% occupancy) showed a dose-dependent occupancy of D₂ receptors with an ED₅₀ values of 5.27 mg/kg (CI = 3.09–7.45) at the 1-h time point. ACR16, on the other hand, showed a dose-dependent occupancy of D₂ receptors over a range of 10 to 240 mg/kg (35–90% occupancy) with an ED₅₀ of 18.99 mg/kg (CI = 11.59–26.39) 1 h after administration. Likewise, haloperidol, over a dose range of 0.025 to 1 mg/kg, showed an occupancy range of 53 to 90%, with an ED₅₀ of 0.02 mg/kg (CI = 0.012–0.028). (–)-OSU6162 did not show catalepsy in the dose range used for the occupancy study, but in case of ACR16 weak catalepsy was observed in one of five animals at a dose of 120 mg/kg. Haloperidol-treated animals showed catalepsy when D₂ occupancy was \geq 0.1 mg/kg (>80% D₂ receptor occupancy; Fig. 1). The results of the time-course experiment are compiled in Table 1, whereas 0.5 mg/kg haloperidol occupied D₂ receptors for a very long time, 30 mg/kg (–)-OSU6162 had the shortest D₂ occupancy time course.




Fig. 1. Striatal D_2 receptor occupancy 1 h after administration of (–)-OSU6162, ACR16, or haloperidol ($n = 5$ for each dose level). The curves were generated using a nonlinear regression equation representing a rectangular hyperbola [$y = ax/(b + x)$] using SigmaPlot software.

TABLE 1

Time course of striatal D_2 receptor occupancy of (–)-OSU6162, ACR16, and haloperidol (percentage of occupancy)

Values are mean \pm S.D. of striatal D_2 receptor occupancy values obtained using at least four rats at each dose level.

Drug	1 h	2 h	4 h	6 h	8 h	24 h
(–)-OSU6162, 30 mg/kg	84.23 \pm 4.84	70.17 \pm 3.54	29 \pm 12.25	<15%	N.D.	N.D.
ACR16, 60 mg/kg	77.9 \pm 2.93	72.49 \pm 8.8	78.4 \pm 19.01	47.58 \pm 4.07	N.D.	N.D.
Haloperidol, 0.5 mg/kg	94.29 \pm 1.45	90.46 \pm 4.45	87.92 \pm 2.17	N.D.	79.73 \pm 2.07	36.93 \pm 13.28

N.D., not determined.


Behavioral Stabilization – Locomotor Activity. All three drugs were effective in decreasing amphetamine-induced hyperlocomotion (Fig. 2). The ED_{50} (95% CI) values for (–)-OSU6162, ACR16, and haloperidol were 44.7 (39–52), 28.2 (20–34), and 0.05 (0.04–0.06) mg/kg, respectively. In the 30-min habituation phase of locomotor activity before administration of amphetamine, haloperidol at doses of 0.1 and 0.5 mg/kg significantly ($p < 0.05$) reduced exploratory locomotor activity by 45 and 67%, respectively, from vehicle controls; ACR16 did so only at the highest tested dose, 60 mg/kg (42% from vehicle controls), whereas (–)-OSU6162 up to a dose of 60 mg/kg did not significantly decrease exploratory locomotor activity compared with pooled vehicle controls in the tested doses, indicating subtle effects of dopamine stabilizers on this behavior [one-way ANOVA, $F(11,63) = 5.48$ followed by two-sided Dunnett's post hoc test]. (–)-OSU6162 and ACR16 were able to significantly increase locomotor counts in rats habituated to the activity cages in similar dose ranges required to decrease amphetamine-induced hyperlocomotion and occupy striatal D_2 receptors (Fig. 2). (–)-OSU6162 at 60 mg/kg increased locomotor activity by 112% from baseline in habituated rats and at the same dose decreased amphetamine-induced locomotor activity by 65%. ACR16, on the other hand, at a dose of 30 mg/kg, increased locomotor activity in habituated rats by 87%, whereas decreasing locomotor activity by 64% in animals stimulated by amphetamine. Thus, over an occupancy range of 37 to 87% (3–60 mg/kg) for (–)-OSU6162 and 35 to 74% (10–60 mg/kg) for ACR16, we observed both inhibitory and stimulatory effects on

locomotion, depending upon the prevailing baseline motor activity. However, in contrast to the dopamine stabilizers, haloperidol [over an occupancy range of 33–78% (0.01–0.1 mg/kg)] potently inhibited psychostimulant activity and induced catalepsy, but it failed to activate habituated animals showing low baseline activity.

Striatal DOPA Levels in Reserpinized Rats. The dopamine agonist apomorphine (0.5 mg/kg) produced an 80% suppression of striatal DOPA accumulation, whereas both (–)-OSU6162 (30 mg/kg) and ACR16 (30 and 60 mg/kg) showed a weak but statistically significant increase of DOPA levels above vehicle-treated controls (Fig. 3).

Conditioned Avoidance Response. Rats were administered (–)-OSU6162 (30, 60, and 100 mg/kg) using a within-subject design. The mean values of inhibition of avoidance based on repeated observations (20, 90, and 240 min after drug injection) of the same five animals are shown in Fig. 4. No escape failures were observed in the three tested doses of (–)-OSU6162. The maximal inhibition of avoidance (49%) was observed 20 min postdrug administration for a dose of 100 mg/kg; hence, the ED_{50} was not determined at any time point. For ACR16, rats were administered doses of 30, 60, and 120 mg/kg, and the values for inhibition of avoidance are shown based on repeated observations (20, 90, and 240 min and 24 h after ACR16 injection) of seven animals (Fig. 4). No escape failures were observed at the dose of 30 mg/kg. At 60 mg/kg, escape failures were observed (1/7 at 20 min, 5/7 at 90 min, 1/7 at 240 min, and all the rats recovered at 24-h test

Amphetamine induced hyperlocomotor activity



Locomotor activity in habituated rats




Fig. 2. Top, effects of (-)-OSU6162 ($n = 5$ for each dose level), ACR16 ($n = 5$ for each dose), and haloperidol ($n = 6$ for each dose level) (milligrams per kilogram) on locomotor activity measured for 1-h duration after amphetamine or saline administration and expressed as mean \pm S.E.M. The drugs or vehicle were administered 30 min before amphetamine or saline challenge. **, $p < 0.001$ [one-way ANOVA, $F(14,76) = 29.97$; post hoc Dunnett's (two-sided) test with respect to pooled amphetamine treatment]. Bottom, effects of (-)-OSU6162 ($n = 5$ for each dose level), ACR16 ($n = 6$ for each dose level), and haloperidol ($n = 6$ for each dose level) on rats habituated to locomotor activity cage environment. Locomotor activity was measured for 1 h after drug injection, and counts are expressed as the mean \pm S.E.M. **, $p < 0.001$; *, $p < 0.05$ [one-way ANOVA, $F(11,67) = 10.63$; post hoc Dunnett's (two-sided) test with respect to pooled vehicle treatment].




Fig. 3. Effect of apomorphine, (-)-OSU6162, and ACR16 ($n = 5$ for each dose for all the drugs) on the dopamine synthesis rate in reserpinized animals in the rat striatum. Values are expressed as percentage of vehicle treated controls (mean \pm S.E.M.). *, $p < 0.01$ [one-way ANOVA, $F(5,24) = 52.81$; post hoc Dunnett's (two-sided) with respect to vehicle treatment].

point). At 120 mg/kg, escape failures were observed (2/7 at 20 min, 6/7 at 90 min, 6/7 at 240 min, and all the rats recovered at 24-h test point). The ED_{50} value at 90 min was determined

to be 39.83 mg/kg (95% CI = 24.5–52.8) using GraphPad software (GraphPad Software Inc., San Diego, CA), which was equivalent to $\sim 65\%$ D_2 receptor occupancy. Haloperidol administered at doses of 0.01, 0.05, and 0.15 mg/kg inhibited CAR and did not induce escape failures at the time points tested (20, 90, and 240 min and 24 h after drug injection) (Fig. 4). Its ED_{50} value at 90 min was 0.019 mg/kg (95% CI = 0.002–0.03), equivalent to $\sim 50\%$ D_2 receptor occupancy determined using GraphPad software.

Fos Immunohistochemistry. All three drugs significantly induced Fos in the nucleus accumbens as well as dorsolateral striatum (Fig. 5). The amount of Fos induced by 30 mg/kg (-)-OSU6162 and 60 mg/kg ACR16 was equivalent to 0.1 mg/kg haloperidol in the nucleus accumbens. However, in spite of significant Fos induction in the nucleus accumbens with (-)-OSU6162, inhibition of CAR was minimal. Fos induction in the dorsolateral striatum of animals treated with (-)-OSU6162 as well as ACR16 did not translate into induction of catalepsy, as was the case for haloperidol. Comparing Fos induction to occupancy percentages, both (-)-OSU6162 and ACR16 significantly induced Fos in the nucleus accumbens as well as in the dorsolateral striatum at doses that exceeded 60% D_2 receptor occupancy.




Fig. 4. Effect of (–)-OSU6162 ($n = 5$), ACR16 ($n = 7$), and haloperidol ($n = 6$) on the performance of conditioned avoidance response in rats after single subcutaneous injection. The animals served as their own controls using a within-subject design. Values of percentage inhibition of avoidance are expressed as mean \pm S.E.M. The avoidance values were analyzed for each drug in a repeated measures analysis of variance with dose (vehicle and three drug doses) as a within-subject factor for each time point separately. If the sphericity assumption was not met, Huynh-Feldt correction was applied and the main effect of dose was significant at least at one time point for all the drugs. Post hoc comparisons were performed using the Bonferroni adjustment for multiple comparisons and the level of significance indicated in the figure is that with respect to vehicle treatment (*, $p < 0.05$).

Fig. 5. Fos expression in the nucleus accumbens and dorsolateral striatum due to drug treatment ($n = 4$ for each dose). Rats were killed 2 h after drug administration, and Fos immunoreactive nuclei expressed as the mean \pm S.E.M. were counted within a $400 \times 400\text{-}\mu\text{m}$ grid in the specified brain regions (*, $p < 0.005$) [one-way ANOVA, $F(13,50) = 41.69$; post hoc Dunnett's (two-sided) test with respect to pooled vehicle control of nucleus accumbens]. #, $p < 0.05$ [one-way ANOVA, $F(13,50) = 19.43$; post hoc Dunnett's (two-sided) test with respect to pooled vehicle treatment of dorsolateral striatum].

Plasma Prolactin Measurements. (–)-OSU6162 and haloperidol showed dose-related prolactin induction (Fig. 6). In the haloperidol group, one rat in the 0.05 mg/kg treatment group was a significant outlier (prolactin value of 140 ng/ml; Grubbs test $Z = 1.71$, $p < 0.05$) and was excluded from the calculations. ACR16 elevated prolactin values minimally and a statistically significant increase was noted only at the dose of 120 mg/kg (Fig. 4). For (–)-OSU6162, the 10-mg/kg dose correlated to a central D_2 receptor occupancy of 64% and

induced significant prolactin levels, whereas for ACR16, the 60-mg/kg dose, which induces a D_2 receptor occupancy of 75%, did not significantly increase plasma prolactin levels.

Discussion

The dopamine stabilizers (–)-OSU6162 and ACR16 show a tone-dependent mixture of stimulatory and depressant effects in locomotor activity models, lending credence to the




Fig. 6. Plasma prolactin levels after (–)-OSU6162, ACR16 or haloperidol treatment measured from plasma samples obtained from the occupancy experiment 1-h after subcutaneous administration (minimum of $n = 4$ for each dose). Values are expressed as mean \pm S.E.M. **, $p < 0.001$; *, $p < 0.01$ [one-way ANOVA, $F(9,49) = 15.04$; post hoc Dunnett's (two-sided) test with respect to the pooled vehicle control].

view that their “dopamine stabilization” properties can be identified and replicated (Fig. 2). Despite their reported low in vitro D_2 receptor affinities the drugs show remarkable D_2 occupancies in vivo in a dose-dependent manner, reaching saturation, consistent with an orderly binding to the dopamine D_2 receptors (Fig. 1). As with other antipsychotics, the drugs did show activity in the CAR model, blocked *d*-amphetamine-induced hyperactivity and induced Fos in limbic regions. However, the effects of these drugs were clearly different from the standard D_2 blocker haloperidol, because catalepsy was not observed in any animal receiving (–)-OSU6162 and in only one animal treated with 120 mg/kg ACR16 for which D_2 receptor occupancies in vivo exceeded 80%. At this degree of striatal D_2 receptor occupancy, both typical and atypical antipsychotics act as full antagonists inducing catalepsy in rodents (Wadenberg et al., 2001; Kapur et al., 2003). Also these behavioral stabilizers had minimal inhibitory effects on the exploratory phase during locomotor experiments in rats that were not habituated to locomotor boxes. Our findings of significant D_2 receptor occupancy with (–)-OSU6162 in vivo are in line with previous findings showing that this compound displaced the dopamine agonist Di-PR-5,6-ADTN in vivo in the rat striatum after subcutaneous dosing (Sonesson et al., 1994) and showed full occupancy of striatal D_2 receptors labeled with raclopride after intravenous dosing to anesthetized monkeys (Ekesbo et al., 1999). The implication of the current findings regarding the potential of these drugs to act as antipsychotics in the clinic as well as limitations and caveats are discussed below.

The high level of in vivo receptor occupancy with relatively modest effects on motor activity made us consider whether these compounds may have intrinsic efficacy (i.e., whether they are partial dopamine agonists). Partial D_2 agonists (e.g., aripiprazole or preclamol) have little or no effect on dopamine turnover, whereas full D_2 antagonists cause a significant increase in dopamine turnover (Semba et al., 1995; Oshiro et al., 1998; Nakai et al., 2003; Jordan et al., 2004). To rule out partial agonism by (–)-OSU6162 and ACR16, we used a dopamine-depleted (reserpinized) animal preparation, which is a particularly sensitive assay for in vivo D_2 agonist effects (Yasuda et al., 1988; Petterson et al., 2002). We observed that apomorphine, a full D_2 receptor agonist, produced nearly 80% suppression of striatal DOPA accumulation in reserpinized rats (Fig. 3). Although Sonesson et al. (1994) reported no

effects of 30 mg/kg (–)-OSU6162 on DOPA accumulation in the striatum of reserpinized rats, we found that both (–)-OSU6162 and ACR16 showed weak but statistically significant increases (10–40%) in DOPA accumulation. The ability of (–)-OSU6162/ACR16 to slightly enhance dopamine synthesis rate above reserpine control levels may indicate the presence of synaptic dopamine. Nevertheless, the present data confirm that the two compounds lack intrinsic activity in dopamine autoreceptors displaying a certain degree of supersensitivity (Hjorth et al., 1988). Dopamine antagonists such as haloperidol are reported not to affect dopamine synthesis rate in reserpinized rats (Johansson et al., 1985). In line with the lack of direct D_2 receptor agonist effects of ACR16 and (–)-OSU6162, these compounds are reported to elevate synthesis, release, and metabolism of dopamine in normal rats, as do other full D_2 receptor antagonists (Sonesson et al., 1994; Petterson et al., 2002). In contrast to dopamine agonists (full or partial), (–)-OSU6162 failed to induce behavioral activation in reserpinized rats (Sonesson et al., 1994) and to induce rotational behavior in 6-hydroxy-dopamine-lesioned rats (Nichols et al., 2002), whereas it blocked rotational behavior induced by dopamine agonists such as quinpirole, apomorphine, or L-DOPA in lesioned rats or monkeys without inducing strong akinesia or dystonia (Ekesbo et al., 1997, 2000; Nichols et al., 2002). Finally, we compared (–)-OSU6162 and ACR16 with haloperidol regarding their effects on plasma prolactin levels. Like haloperidol, (–)-OSU6162 and ACR16 induced dose-dependent increase (>100% over baseline) in prolactin, which further confirms their lack of direct agonist activity at dopamine D_2 receptors (Fig. 6). In summary, our data strongly suggest that both compounds act as full dopamine D_2 receptor antagonists in vivo without evidence for intrinsic agonist activity. However, we cannot rule out the possibility that the difference in the behavioral stabilizing property between these drugs and haloperidol could be due to actions at receptors other than D_2 , although significant in vivo or in vitro affinities for other neurotransmitter receptors have not been found (Sonesson et al., 1994; Petterson et al., 2002).

Both compounds showed clear evidence of antipsychotic-like efficacy in the animal models used. In the CAR assay, which is believed to predict clinical efficacy against positive symptoms (Janssen and Awouters, 1994), ACR16 was clearly more efficacious than (–)-OSU6162 (Fig. 5). This difference

could be explained by a comparable D₂ receptor occupancy of (–)-OSU6162 and ACR16 in spite of higher in vitro affinity exhibited by (–)-OSU6162. In addition, (–)-OSU6162 has a shorter time course of D₂ receptor occupancy and that might explain its low efficacy (Table 1). Elevated prolactin levels because of (–)-OSU6162, compared with ACR16 (Fig. 6) at comparative D₂ occupancy levels, could mean differences in its kinetics of distribution (Kapur et al., 2002). Our D₂ receptor occupancy data for ACR16 are similar to those recently published by Carlsson and Carlsson (2006) in which approximately 60 to 70% displacement of raclopride at doses of 150 μmol/kg (approximately 50 mg/kg) was observed. It should be pointed out that the doses at which the compounds are active in the CAR assay are comparatively high (60–120 mg/kg) compared with clinically used agents (Wadenberg et al., 2001; Kapur et al., 2003). However, initial pilot studies with (–)-OSU6162 (100 mg of daily dose) suggest antipsychotic activity in schizophrenic patients when administered as an add-on to standard treatments (Gefvert et al., 2000; Carlsson et al., 2004). It will certainly be of interest to see whether compounds of this class show antipsychotic activity when given as stand-alone therapies. Additional evidence of antipsychotic-like activity of ACR16 and (–)-OSU6162 include preferential Fos induction in the nucleus accumbens compared with the dorsolateral striatum, indicating a certain limbic selectivity (Fig. 5). The high Fos induction because of (–)-OSU6162 compared with ACR16 (in a similar D₂ occupancy range) remains puzzling and could be due to action at targets unaccounted for in the present study. In addition, both compounds completely reversed *d*-amphetamine-induced hyperactivity without inducing strong hypoactivity or catalepsy. ACR16 and in particular (–)-OSU6162 showed evidence of enhanced plasma prolactin levels 1-h post treatment. Data from time-course studies is needed to reveal whether these effects are sustained and therefore of potential clinical significance. Recently, (–)-OSU6162 has also been shown to reduce apomorphine- and amphetamine-induced behavior in subhuman primates that further substantiates a role for low-affinity dopamine D₂ antagonists in the treatment of psychosis (Brandt-Christensen et al., 2006).

It remains a puzzle as how a D₂ blocker could lead to both an increase as well as a decrease in locomotor activity when haloperidol does not do it. The only other antipsychotic tested in a similar manner that showed a similar profile was remoxipride (Sonesson et al., 1994), which is also a low-affinity D₂ antagonist. It is not clear whether low affinity to D₂ receptor is solely responsible for such actions, but clearly, these compounds showed weak but significant behavioral-activating properties in habituated rats both in our hands as well as in other laboratories (Sonesson et al., 1994; Waters et al., 2002). Recently, a mechanism by which these compounds exert their stabilizing properties based on differences in extrasynaptic versus synaptic dopamine neurotransmission has been proposed (Carlsson and Carlsson, 2006). In addition, a recent study of these compounds showed reversal of (+)-MK-801-induced behavioral impoverishment in mice and social withdrawal in rats (Nilsson et al., 2004; Rung et al., 2005). This suggests that the compounds may possess activity against negative symptoms, including social withdrawal. Anecdotal clinical data generated with (–)-OSU6162 lends support to these findings (Gefvert et al., 2000; Carlsson et al., 2004). As is the case for atypical antipsychotics (–)-OSU6162 and

ACR16 (Sonesson et al., 1997; Waters et al., 2002) were reported to enhance dopamine release in the prefrontal cortex of the rat in vivo, a key factor in improving cognitive deficits seen in schizophrenic patients (Abi-Dargham and Moore, 2003). Together, these findings indicate that the dopamine D₂ receptor stabilizers could be effective in treating positive as well as negative and cognitive symptoms in schizophrenia.

Acknowledgments

We acknowledge Jun Parkes and Doug Hussey (PET Centre, Centre for Addiction and Mental Health, Toronto, ON, Canada) for help and the following co-workers from Eli Lilly Research Laboratories (Indianapolis, IN): Jikesh Shah and Chad Wolfe for the synthesis of ACR16 and (–)-OSU6162 and Julie Falcone and Ken Perry for performing the reserpine experiment.

References

- Abi-Dargham A (2004) Do we still believe in the dopamine hypothesis? New data bring new evidence. *Int J Neuropsychopharmacol* **7** (Suppl 1):S1–S5.
- Abi-Dargham A and Moore H (2003) Prefrontal DA transmission at D₁ receptors and the pathology of schizophrenia. *Neuroscientist* **9**:404–416.
- Barbeau A (1961) Dopamine and basal ganglia diseases. *Arch Neurol* **4**:97–102.
- Brandt-Christensen M, Andersen MB, Fink-Jensen A, Werge T, and Gerlach J (2006) The substituted (S)-3-phenylpiperidine (–)-OSU6162 reduces apomorphine- and amphetamine-induced behaviour in *Cebus apella* monkeys. *J Neural Transm* **113**:11–19.
- Carlsson A and Carlsson ML (2006) A dopaminergic deficit hypothesis of schizophrenia: the path to discovery. *Dialogues Clin Neurosci* **8**:137–142.
- Carlsson ML (2002) Experimental hypoglutamatergia model for cognitive deficits of schizophrenia: effects of classical neuroleptics, 5-HT_{2A} receptor blocking antipsychotics, the ampakine CX516 and the dopaminergic stabiliser ACR16. *Eur Neuropsychopharmacol* **12** (Suppl 3):S167–S168.
- Carlsson ML, Carlsson A, and Nilsson M (2004) Schizophrenia: from dopamine to glutamate and back. *Curr Med Chem* **11**:267–277.
- Deutch AY, Lee MC, and Iadarola MJ (1992) Regionally specific effects of atypical antipsychotic drugs on striatal Fos expression: the nucleus accumbens shell as a locus of antipsychotic action. *Mol Cell Neuroscience* **3**:332–341.
- Ekesbo A, Andren PE, Gunne LM, Sonesson C, and Tedroff J (2000) Motor effects of (–)-OSU6162 in primates with unilateral 6-hydroxydopamine lesions. *Eur J Pharmacol* **389**:193–199.
- Ekesbo A, Andren PE, Gunne LM, and Tedroff J (1997) (–)-OSU 6162 inhibits levodopa-induced dyskinesias in a monkey model of Parkinson's disease. *Neuroreport* **8**:2567–2570.
- Ekesbo A, Torstenson R, Hartvig P, Carlsson A, Sonesson C, Waters N, Tedroff J, and Langstrom B (1999) Effects of the substituted (S)-3-phenylpiperidine (–)-OSU6162 on PET measurements of [¹¹C]SCH23390 and [¹¹C]raclopride binding in primate brains. *Neuropharmacology* **38**:331–338.
- Gefvert O, Lindstrom L, Dahlback O, Sonesson C, and Waters N (2000) (–)-OSU6162 induces a rapid onset of antipsychotic effect after a single dose. A double-blind placebo-controlled pilot study, in *Proceedings of the Annual Meeting Scandinavian Society for Psychopharmacology*; 1997, April 10–13; Copenhagen, Denmark, pp 93–94, Society for Psychopharmacology, Tranekaer, Denmark.
- Hjorth S, Clark D, and Carlsson A (1988) Dopamine (DA) autoreceptor efficacy of 3-PPP enantiomers after short-term synaptic DA deprivation. *Eur J Pharmacol* **152**:207–215.
- Janssen PA and Awouters FH (1994) Is it possible to predict the clinical effects of neuroleptics from animal data? Part V: from haloperidol and pipamperone to risperidone. *Arzneimittelforschung* **44**:269–277.
- Johansson AM, Arvidsson LE, Hacksell U, Nilsson JL, Svensson K, Hjorth S, Clark D, Carlsson A, Sanchez D, Andersson B, et al. (1985) Novel dopamine receptor agonists and antagonists with preferential action on autoreceptors. *J Med Chem* **28**:1049–1053.
- Jordan S, Koprivica V, Dunn R, Tottori K, Kikuchi T, and Altar CA (2004) In vivo effects of aripiprazole on cortical and striatal dopaminergic and serotonergic function. *Eur J Pharmacol* **483**:45–53.
- Kapur S, Langlois X, Vinken P, Megens AA, De Coster R, and Andrews JS (2002) The differential effects of atypical antipsychotics on prolactin elevation are explained by their differential blood-brain disposition: a pharmacological analysis in rats. *J Pharmacol Exp Ther* **302**:1129–1134.
- Kapur S, VanderSpek SC, Brownlee BA, and Nóbrega JN (2003) Antipsychotic dosing in preclinical models is often unrepresentative of the clinical condition: a suggested solution based on in vivo occupancy. *J Pharmacol Exp Ther* **305**:625–631.
- Nakai S, Hirose T, Uwahodo Y, Imaoka T, Okazaki H, Miwa T, Nakai M, Yamada S, Dunn B, Burris KD, et al. (2003) Diminished catalepsy and dopamine metabolism distinguish aripiprazole from haloperidol or risperidone. *Eur J Pharmacol* **472**: 89–97.
- Neu H, Hartvig P, Torstenson R, Fasth KJ, Sonesson C, Waters N, Carlsson A, Tedroff J, and Langstrom B (1997) Synthesis of [¹¹C-methyl](–)-OSU6162, its regional brain distribution and some pharmacological effects of (–)-OSU6162 on the dopaminergic system studied in the rhesus monkey by positron emission tomography. *Nucl Med Biol* **24**:507–511.

- Nichols NF, Cimini MG, Haas JV, Staton BA, Tedroff J, and Svensson KA (2002) PNU-96391A (OSU6162) antagonizes the development of behavioral sensitization induced by dopamine agonists in a rat model for Parkinson's disease. *Neuropharmacology* **43**:817–824.
- Nieoullon A and Coquerel A (2003) Dopamine: a key regulator to adapt action, emotion, motivation and cognition. *Curr Opin Neurol* **16** (Suppl 2):S3–S9.
- Nilsson M, Carlsson A, Markinhuhta KR, Sonesson C, Pettersson F, Gullme M, and Carlsson ML (2004) The dopaminergic stabiliser ACR16 counteracts the behavioural primitivization induced by the NMDA receptor antagonist MK-801 in mice: implications for cognition. *Prog Neuropsychopharmacol Biol Psychiatry* **28**:677–685.
- Oshiro Y, Sato S, Kurahashi N, Tanaka T, Kikuchi T, Tottori K, Uwahodo Y, and Nishi T (1998) Novel antipsychotic agents with dopamine autoreceptor agonist properties: synthesis and pharmacology of 7-[4-(4-phenyl-1-piperazinyl)butoxy]-3,4-dihydro-2(1H)-quinolinone derivatives. *J Med Chem* **41**:658–667.
- Paxinos G and Watson C (1986) *The Rat Brain in Stereotaxic Coordinates*. Academic Press, New York.
- Pettersson F, Gullme M, Jansson AC, Kullingsjo J, Ljung E, Sonniksen K, Thorngren M, Waters N, and Waters S (2002) The development of ACR16. A new class of dopaminergic stabilizers, in *2002 Abstract Viewer/Itinerary Planner: Program No. 894.11*, Society for Neuroscience Annual Meeting; 2002 November 2–7; Orlando, FL, Society of Neuroscience, Washington, DC.
- Robertson GS, Matsumura H, and Fibiger HC (1994) Induction patterns of Fos-like immunoreactivity in the forebrain as predictors of atypical antipsychotic activity. *J Pharmacol Exp Ther* **271**:1058–1066.
- Rung JP, Carlsson A, Markinhuhta KR, and Carlsson ML (2005) The dopaminergic stabilizers (–)-OSU6162 and ACR16 reverse (+)-MK-801-induced social withdrawal in rats. *Prog Neuropsychopharmacol Biol Psychiatry* **29**:833–839.
- Seeman P (1987) Dopamine receptors and the dopamine hypothesis of schizophrenia. *Synapse* **1**:133–152.
- Semba J, Watanabe A, Kito S, and Toru M (1995) Behavioural and neurochemical effects of OPC-14597, a novel antipsychotic drug, on dopaminergic mechanisms in rat brain. *Neuropharmacology* **34**:785–791.
- Sonesson C, Lin CH, Hansson L, Waters N, Svensson K, Carlsson A, Smith MW, and Wikstrom H (1994) Substituted (S)-phenylpiperidines and rigid congeners as preferential dopamine autoreceptor antagonists: synthesis and structure-activity relationships. *J Med Chem* **37**:2735–2753.
- Sonesson C, Tedroff J, Svensson K, Carlsson A, Hansson LO, and Waters N (1997) (–)-OSU6162. A representative from a unique class of functional modulators of dopaminergic systems, in *Proceedings of the Annual Meeting of the Society for Neuroscience*; 1997 October 25–30; New Orleans, LA, p. 531 (Abstract), Society of Neuroscience, Washington, DC.
- Svensson K, Ekman A, Piercy MF, Hoffmann WE, Lum JT, and Carlsson A (1991) Effects of the partial dopamine receptor agonists SDZ 208-911, SDZ 208-912 and terguride on central monoamine receptors. A behavioral, biochemical and electrophysiological study. *Naunyn-Schmiedeberg's Arch Pharmacol* **344**:263–274.
- Svensson K, Waters N, Sonesson C, Wikstrom H, Nicholas NF, and Carlsson A (1993) (–)-DS121, a novel dopamine D₃ and autoreceptor preferring antagonist: effects on locomotor activity in the rat, in *1993 Proceedings of the Annual Meeting for the Society for Neuroscience*; 1993 November 7–12; Washington, DC, Abstract Part 1:39.16, 80, Society for Neuroscience, Washington, DC.
- Tedroff J, Torstenson R, Hartvig P, Sonesson C, Waters N, Carlsson A, Neu H, Fasth KJ and Langstrom B (1998) Effects of the substituted (S)-3-phenylpiperidine (–)-OSU6162 on PET measurements in subhuman primates: evidence for tone-dependent normalization of striatal dopaminergic activity. *Synapse* **28**:280–287.
- Wadenberg ML, Kapur S, Soliman A, Jones C, and Vaccarino F (2000) Dopamine D2 receptor occupancy predicts catalepsy and the suppression of conditioned avoidance response behavior in rats. *Psychopharmacology* **150**:422–429.
- Wadenberg ML, Soliman A, VanderSpek SC, and Kapur S (2001) Dopamine D(2) receptor occupancy is a common mechanism underlying animal models of antipsychotics and their clinical effects. *Neuropsychopharmacology* **25**:633–641.
- Waters N, Carlsson A, Carlsson M, Martin P, Tedroff J, Kullingsjo J, Pettersson F, Sonesson C, and Waters ES (2002) Pharmacology of ACR16, a dopamine stabilizer, in *2002 Abstract Viewer/Itinerary Planner: Program No. 894.13*, for the Society for Neuroscience; 2002 November 2–7; Orlando, FL, Society for Neuroscience, Washington, DC.
- Waters N, Sonesson C, Wikstrom H, Piercy M, Smith M, Carlsson A, and Svensson K (1993) (–)-DS121 a novel dopamine D₃ and autoreceptor preferring antagonist: biochemical and electrophysiological effects, in *1993 Proceedings of the Annual Meeting for the Society for Neuroscience*; 1993 November 7–12; Washington, DC, Abstract Part 1:39.15, 80, Society for Neuroscience, Washington, DC.
- Yasuda Y, Kikuchi T, Suzuki S, Tsutsui M, Yamada K, and Hiyama T (1988) 7-[3-(4-[2,3-Dimethylphenyl]piperazinyl)propoxy]-2(1H)-quinolinone (OPC-4392), a presynaptic dopamine autoreceptor agonist and postsynaptic D₂ receptor antagonist. *Life Sci* **42**:1941–1954.

Address correspondence to: Dr. Shitij Kapur, Centre for Addiction and Mental Health, ARF Site, 33 Russell St., Toronto, ON, Canada M5S 2S1. E-mail: shitij_kapur@camh.net
